

Charge ordering in quasi-one-dimensional systems with frustrating interactions

Siddhartha Lal^{1,*} and Mukul S. Laad^{2,†}

¹*Institut für Theoretische Physik, Universität zu Köln, Zùlpicher Straße 77, D-50937 Köln, Germany*

²*Department of Physics, Loughborough University, LE11 3TU, UK*

Motivated by the co-existing charge and spin order found in strongly correlated ladder systems, we study an effective pseudospin model on a coupled two-leg ladder. A bosonisation analysis yields a rich phase diagram showing Wigner/Peierls charge order and Neel/dimer spin order. In a broad parameter regime, the orbital antiferromagnetic phase is found to be stable. An intermediate gapless phase of finite width is found to lie in between two charge-ordered gapped phases. Our work is potentially relevant for a unified description of a class of strongly correlated, quarter-filled ladder systems.

PACS numbers: 71.30.+h, 71.10.Pm

Strongly correlated ladder systems are fascinating candidates for studying the interplay of spin and charge ordering, and their combined influence on the emergence of novel ground states [1]. Several recent studies provide experimental realisations of such systems, exhibiting diverse phenomena like charge order (CO), antiferromagnetism (AF) and unconventional superconductivity (uSC) as functions of suitable control parameters [2, 3]. The existence of several non-perturbative theoretical techniques in one dimension have also resulted in the study of the emergence of exotic phases from instabilities of the high- T Luttinger liquid [4]. Given that these systems are Mott insulators, longer-range Coulomb interactions are relevant in understanding CO/AF/uSC phases. In spite of some recent work [5], a detailed understanding of the effects of longer-range interactions in quasi-1D systems is a largely unexplored problem. Further, attention has mostly focussed on studies of models at 1/2-filling [4].

Here, we study an effective pseudospin model describing charge degrees of freedom of a one-dimensional 1/4-filled electronic system. Such an effective model can be derived from an extended Hubbard model (with longer-range interactions) in a number of physically relevant cases [6, 7, 8]. It has also been suggested [9] that this may be a relevant starting point for the study of charge-order in organics. While the weak coupling limit of the underlying extended Hubbard model has been recently studied [10, 11], we note that the real systems under consideration are generically in the strong coupling regime of the model [6, 7, 8]. To the best of our knowledge, this regime has not been studied in sufficient detail.

We start with the effective pseudospin Hamiltonian

$$H^{chain} = - \sum_j [2t \tau_j^x + V \tau_j^z \tau_{j+1}^z] \quad (1)$$

where $2t$ is the transverse field, V is the n.n interaction pseudospin coupling. Here, the τ represent charge degrees of freedom in an effective model derived from a more basic electronic Hamiltonian for a 1/4-filled system [6, 7, 8]. This is the 1D Ising model in a transverse field.

Rotating the pseudospin axis $\tau^x \rightarrow \tau^z$, $\tau^z \rightarrow -\tau^x$ and introducing a bond-fermion repulsion $U_\perp \sum_{i,a,b \neq a} (n_{i,a} - n_{i+1,a})(n_{i,b} - n_{i+1,b})$ as well as a bond-fermion transfer term $t_\perp \sum_{i,a,b \neq a} (c_{i,a,\uparrow}^\dagger c_{i+1,a,\uparrow} c_{i,b,\downarrow}^\dagger c_{i+1,b,\downarrow} + \text{h.c.})$ between two such systems described by the indices (a, b) [6, 7, 8], we have the effective Hamiltonian for the charge sector of the coupled system in terms of a pseudospin ladder model

$$H = - \sum_{j,a} [2t \tau_{j,a}^z + V \tau_{j,a}^x \tau_{j+1,a}^x] - \sum_{j,a,b \neq a} [U_\perp \tau_{j,a}^z \tau_{j,b}^z + t_\perp (\tau_{j,a}^x \tau_{j,b}^x + \tau_{j,a}^y \tau_{j,b}^y)], \quad (2)$$

where $a, b = 1, 2$ is the chain index. Having explored the case of $U_\perp \gg V$ in an earlier work [12], we will explore the scenario of $U_\perp \ll V$ below. In the spin sector, coupling between the two systems leads to a $S = 1/2$ Heisenberg ladder-type model, and has been studied extensively by several authors [4, 13]. Interchain spin coupling turns out to be relevant, opening up a spin gap concomitantly with generating AF/dimer ordered ground states with spontaneously broken SU(2)/translation symmetries.

We now analyse the charge sector. To begin, we introduce (bond-)fermion operators via a Jordan-Wigner transformation. In terms of these fermions, upon denoting the chains as $a = \uparrow, \downarrow$, we find an effective Hamiltonian for the 1D Hubbard model with an equal-spin pairing term and an on-site spin-flip term

$$H = - \frac{\tilde{t}}{2} \sum_{j,a} (\psi_{j,a}^\dagger \psi_{j+1,a} + \text{h.c.}) - \frac{t_\perp}{2} \sum_j (\psi_{j,a}^\dagger \psi_{j,b} + \text{h.c.}) + \tilde{U} \sum_j n_{j,\uparrow} n_{j,\downarrow} + \tilde{V} \sum_{j,a} (\psi_{j,a}^\dagger \psi_{j+1,a}^\dagger + \text{h.c.}) + \mu \sum_{j,a} n_{j,a} \quad (3)$$

where $\tilde{t}$ and $t_\perp$ are the in-chain and inter-chain hopping parameters respectively, $\mu = -2t$ the chemical potential, $\tilde{U} = -U_\perp$ the on-site (Hubbard) interaction coupling and $\tilde{V} = V/4$ the pairing strength. Note that while we treat the parameters $\tilde{t}$ and $\tilde{V}$ as independent parameters for the sake of generality, $\tilde{t} = \tilde{V} = V$ in our original model

(2). Bosonising in the usual way (in terms of the usual charge $\rho = \uparrow + \downarrow$ and spin $\sigma = \uparrow - \downarrow$ variables), we obtain the effective low-energy bosonic Hamiltonian

$$\begin{aligned}
H = & \frac{1}{2\pi} \int dx [v_\rho K_\rho (\pi \Pi_\rho(x))^2 + \frac{v_\rho}{K_\rho} (\partial_x \phi_\rho(x))^2] \\
& + \frac{1}{2\pi} \int dx [v_\sigma K_\sigma (\pi \Pi_\sigma(x))^2 + \frac{v_\sigma}{K_\sigma} (\partial_x \phi_\sigma(x))^2] \\
& + \frac{U_\sigma}{2\pi\alpha} \int dx \cos(\sqrt{8}\phi_\sigma) + \frac{U_\rho}{2\pi\alpha} \int dx \cos(\sqrt{8}\phi_\rho) \\
& + \frac{\tilde{V}}{2\pi\alpha} \int dx \cos(\sqrt{2}\theta_\rho) \cos(\sqrt{2}\theta_\sigma) - \frac{\sqrt{2}\mu}{\pi} \int dx \partial_x \phi_\rho(x) \\
& + \frac{t_\perp}{\pi\alpha} \int dx \cos(\sqrt{2}\phi_\sigma) \cos(\sqrt{2}\theta_\sigma) + \frac{V_1}{\pi\alpha} \int dx \cos(\sqrt{8}\theta_\sigma) \\
& + \frac{V_2}{\pi\alpha} \int dx \cos(\sqrt{2}\phi_\sigma) \cos(\sqrt{2}\theta_\rho) + \frac{V_3}{\pi\alpha} \int dx \cos(\sqrt{8}\theta_\rho) \quad (4)
\end{aligned}$$

where $\Pi_\rho = \frac{1}{\pi} \partial_x \theta_\rho$, $\Pi_\sigma = \frac{1}{\pi} \partial_x \theta_\sigma$, $v_\rho K_\rho = v_F = v_\sigma K_\sigma$, $v_\rho/K_\rho = v_F(1 + \frac{\tilde{U}}{\pi v_F})$ and $v_\sigma/K_\sigma = v_F(1 - \frac{\tilde{U}}{\pi v_F})$. Among the various cosine potentials, we have the usual spin-flip backscattering $\cos(\sqrt{8}\phi_\sigma)$ and Umklapp $\cos(\sqrt{8}\phi_\rho)$ terms as well as the triplet superconducting $\cos(\sqrt{2}\theta_\rho) \cos(\sqrt{2}\theta_\sigma)$ term. The chemical potential term can be absorbed by performing the shift $\phi_\rho \rightarrow \phi_\rho + \frac{\sqrt{2}K_\rho\mu}{v_\rho}x$. The cosine potentials with couplings V_1 , V_2 and V_3 are generated under RG by the $t_\perp$ and $\tilde{V}$ terms. We find the RG equations for the various couplings to second-order as

$$\begin{aligned}
\frac{dU_\rho}{dl} &= (2 - 2K_\rho)U_\rho \\
\frac{dU_\sigma}{dl} &= (2 - 2K_\sigma)U_\sigma - (\frac{1}{K_\sigma} - K_\sigma)t_\perp^2 \\
\frac{d\tilde{V}}{dl} &= (2 - \frac{1}{2}(\frac{1}{K_\sigma} + \frac{1}{K_\rho}))\tilde{V} - K_\sigma t_\perp V_2 \\
\frac{dt_\perp}{dl} &= (2 - \frac{1}{2}(K_\sigma + \frac{1}{K_\sigma}))t_\perp - \frac{\tilde{V}V_2}{K_\rho} - (K_\sigma U_\sigma + \frac{V_1}{K_\sigma})2t_\perp \\
\frac{dV_1}{dl} &= (2 - \frac{2}{K_\sigma})V_1 + (\frac{1}{K_\sigma} - K_\sigma)t_\perp^2 \\
\frac{dV_2}{dl} &= (2 - \frac{1}{2}(K_\sigma + \frac{1}{K_\rho}))V_2 - \frac{t_\perp \tilde{V}}{K_\sigma} \\
\frac{dV_3}{dl} &= (2 - \frac{2}{K_\rho})V_3. \quad (5)
\end{aligned}$$

The RG equations for the two interaction parameters (K_ρ, K_σ) as well as the parameter $\delta = K_\rho\mu/v_\rho$ are found to be

$$\begin{aligned}
\frac{dK_\sigma}{dl} &= -K_\sigma^2 U_\sigma^2 + V_1^2 \\
\frac{dK_\rho}{dl} &= -K_\rho^2 U_\rho^2 J_0(\delta(l)\alpha) \\
\frac{d\delta}{dl} &= \delta(l) - U_\rho^2 J_1(\delta(l)\alpha), \quad (6)
\end{aligned}$$

where $\delta(l) = \delta e^l$, α is a short-distance cut-off like the lattice spacing and $J_0(x), J_1(x)$ are Bessel functions [4].

For repulsive interactions ($U_\perp > 0$) between the bond-fermions, the σ sector is massless and K_σ flows under RG to the fixed point value $K_\sigma^* \gtrsim 1$ and $1/2 \leq K_\rho \leq 1$. At 1/2-filling (for the bond-fermions), the couplings U_ρ , $\tilde{V}$, $t_\perp$, V_1 and V_2 are all relevant while U_σ and V_3 are irrelevant. The competition to reach strong-coupling first is, however, mainly between U_ρ , $t_\perp$ and $\tilde{V}$. We show below the phase diagram as derived from this analysis.




FIG. 1: The RG phase diagram in the (K_σ, K_ρ) plane for repulsive interchain interactions ($U_\perp < 0$). The three regions $K_\rho < (1/4(1/K_\sigma + 1/K_\rho), 1/4(K_\sigma + 1/K_\sigma))$, $1/K_\sigma > K_\rho > 1/4(1/K_\sigma + 1/K_\rho)$ and $K_\rho > (1/K_\sigma, 1/4(K_\sigma + 1/K_\sigma))$ give the values of (K_σ, K_ρ) for which the couplings U_ρ , $t_\perp$ and $\tilde{V}$ respectively are the fastest to grow under RG.

In the phase diagram in Fig.(1), the three lines with intercepts at $(K_\sigma = 1, K_\rho = 1)$, $(K_\sigma = 1, K_\rho = 0.64)$ and $(K_\sigma = 1, K_\rho = 1/2)$ are the relations $K_\sigma = 1/K_\rho$, $K_\rho = 1/4(1/K_\sigma + 1/K_\rho)$ and $K_\rho = 1/4(K_\sigma + 1/K_\sigma)$ respectively. The regions $K_\rho < (1/4(1/K_\sigma + 1/K_\rho), 1/4(K_\sigma + 1/K_\sigma))$, $1/K_\sigma > K_\rho > 1/4(1/K_\sigma + 1/K_\rho)$ and $K_\rho > (1/K_\sigma, 1/4(K_\sigma + 1/K_\sigma))$ signify the values of K_ρ and K_σ for which U_ρ (in-chain Wigner charge-ordered Mott insulator), $t_\perp$ and $\tilde{V}$ (in-chain Peierls charge-ordered Mott insulator of preformed bond-fermion pairs) respectively are the fastest to reach strong-coupling. The RG equations for the coupling U_ρ , the interaction parameter K_ρ and the incommensuration parameter δ are familiar from the literature on commensurate-incommensurate transitions [14]. For temperatures $T \gg K_\rho\mu$, the finite chemical potential is unable to quench the Umklapp scattering processes, allowing for the growth of U_ρ to strong-coupling. For $T \ll K_\rho\mu$, the finite chemical potential cuts off the RG flow of U_ρ , freezing the Umklapp scattering processes.

For the case of $t_\perp$ being the most relevant coupling, we find our RG equations to be a non-trivial generalisation of those derived in [15] for the model of two coupled spinless fermion chains without the intrachain

$\tilde{V}$ pairing term. The resulting picture then describes strong interchain two-particle correlations between bond-fermions sharing a rung. Following the analysis outlined in [13, 15], we conclude that this phase is a novel insulating phase characterised by a mass gap and delocalised bond-fermions on rungs, resembling the orbital antiferromagnetic ground state found in the spinless two-chain problem. This matches our finding of an orbital antiferromagnetic ground state in the strongly-coupled ladder with dominant antiferromagnetic rung-couplings in an earlier work [12]. Away from 1/2-filling (for the bond-fermions), the competition is mainly between $t_\perp$ and $\tilde{V}$. For $\tilde{V}$ reaching strong-coupling ahead of $t_\perp$, the system is in a channel triplet-spin singlet superconducting phase with mobile intra-chain hole pairs; for $t_\perp$ reaching strong-coupling ahead of $\tilde{V}$, we are currently unable to describe in more detail the dominant instability away from the orbital antiferromagnetism like insulating phase away from quarter-filling.

Interestingly, while studies of ladder models have shown a plethora of charge and spin-ordered gapped phases [4, 10, 11], there remains the intriguing possibility that some of these gapped phases may themselves be separated from one another by non-trivial gapless phases of finite width [13]. In what follows, we provide an explicit realisation of this scenario. The RG equations (5)-(6) reveal the existence of a non-trivial fixed point (FP) for any value of (K_ρ, K_σ) lying in the ranges $1/2 < K_\rho < 1$, $1 < K_\sigma < 2 + \sqrt{3}$ and which is perturbatively accessible from the trivial weak-coupling FP. We note that a similar non-trivial fixed point was found in a study of the anisotropic Heisenberg spin-1/2 chain in a magnetic field [16], where the authors derived a set of RG equations which were very similar to those found in [15]. Here, the non-trivial FP is given by

$$t_\perp^* = \sqrt{ab}, \quad V_1^* = \frac{K_\sigma^* + 1}{2} t_\perp^{*2}, \quad U_\sigma^* = \frac{V_1^*}{2K_\sigma^*}$$

$$\tilde{V}^* = \sqrt{aK_\rho^*(cK_\sigma^* - (K_\sigma^* + 1)^2 ab)}, \quad V_2^* = \sqrt{\frac{b}{a} \frac{\tilde{V}^*}{K_\sigma^*}} \quad (7)$$

where $a = 2 - (K_\sigma^* + 1/K_\rho^*)/2$, $b = 2 - (1/K_\sigma^* + 1/K_\rho^*)/2$ and $c = 2 - (K_\sigma^* + 1/K_\sigma^*)/2$. Further, we can safely make the approximation of the renormalisations of K_ρ and K_σ being small at this non-trivial FP [16]. The system is gapless at this non-trivial FP as well as at points which flow to it. The trivial FP has 6 unstable directions ($U_\rho, \tilde{V}, t_\perp, V_1, V_2$ and δ), 2 stable directions (U_σ and V_3) and 2 marginal directions (K_ρ and K_σ). The non-trivial FP has 5 unstable directions, 3 stable directions and two marginal directions. Apart from the stable direction given by the V_3 , the presence of the other two stable directions at the non-trivial FP indicates the existence of a two-dimensional surface of gapless theories in the five-dimensional $(U_\sigma, \tilde{V}, t_\perp, V_1, V_2)$ coupling space. This gapless phase is the analog of the ‘‘Floating Phase’’

found in the phase diagram of the 1D axial next nearest neighbour Ising model [16]. We present in Fig.(2) below a RG flow phase diagram which is projected onto the $(V_1, t_\perp)$ plane (a similar RG flow diagram is found for the case of the anisotropic Heisenberg model in a magnetic field h [16] in the (a, h) plane, where a is the anisotropy parameter).




FIG. 2: The RG phase diagram in the $(V_1, t_\perp)$ plane. The thick line characterises the set of points which flow to the intermediate fixed point $(V_1^*, t_\perp^*)$ shown by the filled circle. The thin lines show all RG flows which flow towards strong-coupling in the two phases **I** and **II**, characterised by the relevant couplings $\tilde{V}$ and $t_\perp$ respectively.

The regions **I** and **II** characterise all RG flows which do not flow to the intermediate fixed point at $(V_1^*, t_\perp^*)$. In region **I**, V_1 flows to strong-coupling while $t_\perp$ decays; for $1/2 < K_\rho < 1$ and $K_\sigma > 1$, we know from the above discussion that in this region, the coupling $\tilde{V}$ will reach strong-coupling first. In region **II**, both $t_\perp$ and V_1 grow under RG, with the coupling $t_\perp$ being the first to reach strong-coupling. Thus, the RG trajectory leading to the intermediate fixed point represents a gapless phase separating the two gapped, charge-ordered phases **I** and **II** characterised by the relevant couplings $\tilde{V}$ and $t_\perp$ respectively.

For attractive interactions ($U_\perp < 0$) between the bond-fermions, we can carry out a similar analysis. In this case, we can see that $K_\rho > 1$ while $K_\sigma < 1$. Then, from the RG equations given above, we can see that the Umklapp coupling U_ρ and V_1 are irrelevant while the couplings $t_\perp, U_\sigma, \tilde{V}, V_2$ and V_3 are relevant. The competition to reach strong-coupling first is, however, mainly between $U_\sigma, t_\perp$ and $\tilde{V}$. We show below the phase diagram at 1/2-filling for the bond-fermions as derived from this analysis.

In the phase diagram in Fig.(3), the three lines with intercepts at $(K_\rho = 1, K_\sigma = 1)$, $(K_\rho = 1, K_\sigma = 0.64)$ and $(K_\rho = 1, K_\sigma = 1/\sqrt{3})$ are the relations $K_\sigma = 1/K_\rho$, $K_\sigma = 1/4(1/K_\sigma + 1/K_\rho)$ and $K_\sigma = 1/\sqrt{3}$ respectively.




FIG. 3: The RG phase diagram in the (K_ρ, K_σ) plane for attractive interchain interactions ($U_\perp > 0$). The three regions $K_\sigma < (1/4(1/K_\sigma + 1/K_\rho), 1/\sqrt{3})$, $1/K_\rho > K_\sigma > 1/\sqrt{3}$ and $K_\sigma > (1/K_\rho, 1/4(1/K_\rho + 1/K_\sigma))$ give the values of (K_σ, K_ρ) for which the couplings U_σ , $t_\perp$ and $\tilde{V}$ respectively are the fastest to grow under RG.

The regions $K_\sigma < (1/4(1/K_\sigma + 1/K_\rho), 1/\sqrt{3})$, $1/K_\rho > K_\sigma > 1/\sqrt{3}$ and $K_\sigma > (1/K_\rho, 1/4(1/K_\rho + 1/K_\sigma))$ signify the values of K_ρ and K_σ for which U_σ (rung-dimer insulator with in-chain Wigner charge-ordering), $t_\perp$ and $\tilde{V}$ (insulator with in-chain dimers and Peierls charge-ordering) respectively are the fastest to reach strong-coupling. This matches our finding of a ground state with in-chain Wigner charge order and rung-dimers in the strongly-coupled ladder with large ferromagnetic rung-couplings in an earlier work [12]. Away from 1/2-filling (for the bond-fermions), depending on which of the three couplings $t_\perp$, $\tilde{V}$ and U_σ is the first to reach strong-coupling, the system exists either as a superconductor with intra-chain hole pairs ($\tilde{V}$) or a superconductor with rung-singlet hole pairs (U_σ) or a phase reached by following the dominant instability away from the orbital antiferromagnetism like insulating phase ($t_\perp$) but which we are currently unable to describe in greater detail.

To conclude, we have studied a model of strongly correlated coupled quasi-1D systems at 1/4-filling using an effective pseudospin ladder model with $V > t, t_\perp, U_\perp$. Using a bosonisation analysis, we find two different types of charge/spin ordered ground states at 1/4-filling. Transverse bond-fermion hopping is found to stabilise a new, gapped (insulating) phase characterised by interchain two-particle coherence of a type resembling orbital antiferromagnetism [13, 15]. The spin fluctuations are described by a $S = 1/2$ Heisenberg ladder-type model for all cases studied here: the spin excitations are always

massive. Away from this filling, either intra- or inter-chain superconductivity in a gapped spin background is found to be the stable ground state. We also find the existence of an intermediate gapless phase lying in between two gapped, charge-ordered phases (characterised by the relevant couplings $\tilde{V}$ and $t_\perp$ respectively) in the RG phase diagram of our model. Our analysis is especially relevant to ladder systems like $Sr_{14-x}Ca_xCu_{24}O_{41}$ and $\alpha - NaV_2O_5$ or $\beta - Na_{0.33}V_2O_5$ (a superconductor) which exhibit charge/spin long range order at $x = 0$ and superconductivity beyond under pressure and/or doping [2, 3].

SL and MSL thank the DFG (Germany) and EPSRC (UK) respectively for financial support.

* Electronic address: sanjayl@thp.uni-koeln.de

† Electronic address: M.S.Laad@lboro.ac.uk

- [1] S. Sachdev, *Quantum Phase Transitions* (Cambridge Univ. Press, Cambridge, 1999) and references therein.
- [2] E. M. McCarron III *et al.*, Mater. Res. Bull. **23**, 1355 (1988); M. Uehara *et al.*, J. Phys. Soc. Jpn. **65**, 2764 (1996).
- [3] T. Yamauchi, Y. Ueda and N. Mori, Phys. Rev. Lett. **89**, 057002 (2002).
- [4] Thierry Giamarchi, *Quantum Physics in One Dimension* (Oxford Univ. Press, Oxford, 2004) and references therein.
- [5] S. Capponi, D. Poilblanc and T. Giamarchi, Phys. Rev. B **61**, 13410 (2000).
- [6] P. Thalmeier and P. Fulde, Europhys. Lett. **44**, 242 (1998).
- [7] M. V. Mostovoy and D. I. Khomskii, Solid State Commun. **113**, 159 (2000).
- [8] M. Aichhorn, P. Horsch, W. von der Linder and M. Cuoco, Phys. Rev. B **65**, 201101 (2002).
- [9] V. J. Emery and C. Noguera, Phys. Rev. Lett. **60**, 631 (1988).
- [10] P. Donohue, M. Tsuchiizu, T. Giamarchi and Y. Suzumura, Phys. Rev. B **63**, 045121 (2001).
- [11] E. Orignac and R. Citro, Eur. Phys. J. B **33**, 419 (2003).
- [12] M. S. Laad, S. Lal, cond-mat/0510016.
- [13] A. O. Gogolin, A. A. Nersesyan and A. M. Tsvelik, *Bosonization and Strongly Correlated Systems* (Cambridge Univ. Press, Cambridge, 1998) and references therein.
- [14] G. I. Japaridze and A. A. Nersesyan, Sov. Phys. JETP Lett. **27**, 334 (1978); V. L. Pokrovsky and A. Talapov, Phys. Rev. Lett. **42**, 65 (1979).
- [15] F. V. Kusmartsev, A. Luther and A. A. Nersesyan, JETP Lett. **55**, 692 (1992).
- [16] A. Dutta and D. Sen, Phys. Rev. B **67**, 094435 (2003).